ACL INJURY PREVENTION

Alentorn-Geli E, Mendiguchia J, Samuelsson K, Musahl V, Karlsson J, Cugat R, Myer GD. Prevention of non-contact anterior cruciate ligament injuries in sports. Part II: systematic review of the effectiveness of prevention programmes in male athletes. *Knee Surg Traum Arth*, 2014;22(1):16-25.

DiStefano LJ, Blackburn JT, Marshall SW, Guskiewicz KM, Garrett WE, Padua DA. Effects of an age-specific anterior cruciate ligament injury prevention program on lower extremity biomechanics in children. *Am J Sports Med*, 2011;39(5):949-957.

DiStefano LJ, DiStefano MJ, Frank BS, Clark MA, Padua DA. Comparison of integrated and isolated training on performance measures and neuromuscular control. *J Str Cond Res*, 2013;27(4):1083-1090.

DiStefano LJ, Padua DA, Blackburn JT, Garrett WE, Guskiewicz KM, Marshall SW. Integrated injury prevention program improves balance and vertical jump height in children. *J Str Cond Res*, 2010;24(2):332-342.

DiStefano LJ, Padua DA, DiStefano MJ, Marshall SW. Influence of age, sex, technique, and exercise program on movement patterns after an anterior cruciate ligament injury prevention program in youth soccer players. *Am J Sports Med*, 2009;37(3):495-505.

Dwyer MK, Boudreau SN, Mattacola CG, Uhl TL, Lattermann C. Comparison of lower extremity kinematics and hip muscle activation during rehabilitation tasks between sexes. *J Ath Train*, 2010;45(2):181-190.

Gagnier JJ Morgenstern H, Chess L. Interventions designed to prevent anterior cruciate ligament injuries in adolescents and adults: A systematic review and meta-analysis. *Am J Sports Med*, 2013;41(8):1952-1962.

Hagglund M, Atroshi I, Wagner P, Walden M. Superior compliance with a neuromuscular training programme is associated with fewer ACL injuries and fewer acute knee injuries in female adolescent football players: secondary analysis of an RCT. *Br J Sports Med*, 2013;47:974-979.

Hollman JH, Galardi CM, Lin IH, Voth BC, Whitmarsh CL. Frontal and transverse plane hip kinematics and gluteus maximus recruitment correlate with frontal plane kinematics during single-leg squat tests in women. *Clin Biomech*, 2014;29(4):468-474.

Joy EA, Taylor JR, Novak MA, Chen M, Fink BP, Porucznik CA. Factors influencing the implementation of anterior cruciate ligament injury prevention strategies by girls soccer coaches. *J Str Cond Res*, 2013;27(8):2263-2269.

Lauersen JB, Bertelsen DM, Andersen LB. The effectiveness of exercise interventions to prevent sports injuries: a systematic review and meta-analysis of randomised controlled trials. *Br J Sports Med*, 2014;48:871-877.

Mauntel TC, Frank BS, Begalle RL, Blackburn JT, Padua DA. Kinematic differences between those with and without medial knee displacement during a single-leg squat. *J Appl Biomech*, 2014;30(6):707-712.

McCurdy K, O'Kelley E, Kutz M, Langford G, Ernest J, Torres M. Comparison of lower extremity EMG between the 2-leg squat and modified single-leg squat in female athletes. *J Sport Rehab*, 2010;19(1):57-70

Myklebust G, Skjolberg A, Bahr R. ACL injury incidence in female handball 10 years after the Norwegian ACL prevention study: important lessons learned. *Br J Sports Med*, 2013;47:476-479.

Nguyen, A, Schultz, SJ, Schmitz RJ, Luecht RM, Perrin DH. A preliminary multifactorial approach describing the relationships among lower extremity alignment, hip muscle activation, and lower extremity joint excursion. *J Ath Train*, 2011;46(3):246-256.

Padua DA, DiStefano LJ, Marshall SW, Beutler AI, de la Motte SJ, DiStefano MJ. Retention of movement pattern changes after a lower extremity injury prevention program is affected by program duration. *Am J Sports Med*, 2012;40(2):300-306.

Pappas E, Nightengale EJ, Simic M, Ford KR, Hewett TE, Myer GD. Do exercises used in injury prevention programmes modify cutting task biomechanics? A systematic review with meta-analysis. *Br J Sports Med*, 2014;ePub ahead of print.

Pollard DC, Stearns, KM, Hayes AT, Heiderscheit BC. Altered lower extremity movement variability in female soccer players during sidestep cutting after anterior cruciate ligament reconstruction. *Am J Sports Med*, 2014;ePub ahead of print Dec 2014

Sadoghi P, von Keudell A, Vavken P. Effectiveness of anterior cruciate ligament injury prevention training programs. *J Bone Joint Surg*, 2012;94:769-776.

Schmitt LC, Paterno MV, Ford KR, Myer GD, Hewett TE. Strength asymmetry and landing mechanics at return to sport after ACL reconstruction. *Med Sci Sports Exerc*, 2014;ePub ahead of print Nov 2014.

Steffen K, Meeuwisse WH, Romiti M, Kang J. McKay C, Bizzini M, Dvorak J, Finch C, Myklebust G, Emery CA. Evaluation of how different implementation strategies of an injury prevention proramme (FIFA 11+) impact team adherence and injury risk in Canadian female football players: a cluster-randomized trial. *Br J Sports Med*, 2013;47:480-487.

Sugimoto D, Myer GD, Barber Foss KD, Hewett TE. Dosage effects of neuromuscular training intervention to reduce anterior cruciate ligament injuries in female athletes: Meta- and sub-group analyses. *Sports Med*, 2014;44(4):55`-562.

Taylor JB, Waxman JP, Richter SJ, Shultz SJ. Evaluation of the effectiveness of anterior cruciate ligament injury prevention programme training components: a systematic review and meta-analysis. *Br J Sports Med*, 2013:1-10. Published online first Aug 6 2013.